



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The equation may be written

$$\frac{1}{y_{x+1}} - \frac{1}{y_x} = \frac{y_{x+1}}{xy_x}.$$

Hence

$$\frac{1}{y_x} - \frac{1}{y_1} = \sum_1^{x-1} \frac{1}{x} \cdot \frac{y_{x+1}}{y_x} = \sum_1^{x-1} \frac{1}{x} \left[1 - \frac{y_x - y_{x+1}}{y_x} \right] = \sum_1^{x-1} \frac{1}{x} - \sum_1^{x-1} \frac{1}{x^2} \cdot \frac{y_{x+1}^2}{y_x}.$$

Now

$$\sum_1^{x-1} \frac{1}{x} = \log x + r,$$

where r approaches Euler's constant C when x becomes infinite (Goursat-Hedrick, vol. 1, p. 103, Ex. 1); so that, if we substitute and divide by $\log x$, we have

$$\frac{1}{y_x \log x} - 1 = \frac{\frac{1}{y_1} + r - \sum_1^{x-1} \frac{1}{x^2} \cdot \frac{y_{x+1}^2}{y_x}}{\log x}.$$

But from (2) $y_{x+1} < y_x < y_1$, $y_{x+1}^2 < y_1 y_x$. Therefore

$$\sum_1^{x-1} \frac{1}{x^2} \cdot \frac{y_{x+1}^2}{y_x} < y_1 \sum_1^{x-1} \frac{1}{x^2} < \frac{\pi^2 y_1}{6}.$$

The numerator of the fraction remains numerically less than a fixed number when x becomes infinite and we have

$$\lim y_x \log x = 1.$$

It is to be noticed that the above method can be immediately extended to

$$y_x = f(x)y_{x+1}^2 + y_{x+1},$$

for certain functions $f(x)$, where $\sum_1^\infty [f(x)]^2$ is a convergent series.

NOTE. From (1) we obtain

$$\frac{y_x}{y_{x+1}} - 1 = \frac{y_{x+1}}{x}, \quad \text{and} \quad x \left(\frac{1}{y_{x+1}} - \frac{1}{y_x} \right) = \frac{y_{x+1}}{y_x}. \quad (3)$$

When x becomes infinite, the first equation in (3) together with (2) shows that y_x/y_{x+1} approaches unity. Hence the second equation gives

$$\lim_{x \rightarrow \infty} x \left(\frac{1}{y_{x+1}} - \frac{1}{y_x} \right) = 1.$$

It is easily seen that the limit is the same if we replace x by $x+1$ and when this is done it follows that

$$\lim_{x \rightarrow \infty} \frac{1}{\log(x+1)y_{x+1}} = 1,$$

by use of the theorem on page 108, § 162, E. Cesàro, *Elementares Lehrbuch der algebraischen Analysis* . . ., Leipzig, 1904. The desired result easily follows from the above.

2846 [1920, 326].

Find the entire volume within the surface $x^{1/2} + y^{1/2} + z^{1/2} = a^{1/2}$. (W. A. Granville, *Elements of Differential and Integral Calculus*, revised ed., 1911, p. 420.)

This equation, rationalized, is the equation of Steiner's quartic surface, every tangent plane to which cuts it in two conics. (Cf. Salmon-Rogers, *Analytic Geometry of Three Dimensions*, 5th ed., vol. 2, 1915, pp. 171, 201, 207, 213f. Also C. M. Jessop, *Quartic Surfaces* 1916, chapter 7.)

I. SOLUTION BY L. A. EASTBURN, North Arizona Normal School, Flagstaff, Ariz.

The required volume inclosed by the surface is

$$v = \int_0^a \int_0^{(a^{1/2}-x^{1/2})^2} \int_0^{a^{1/2}-x^{1/2}-y^{1/2}} dz dy dx,$$

$$\begin{aligned}
&= \int_0^a \int_0^{(a^{1/2}-x^{1/2})^2} [(a^{1/2}-x^{1/2})^2 - 2(a^{1/2}-x^{1/2})y^{1/2} + y] dy dx, \\
&= 1/6 \int_0^a [a^2 - 4a^{3/2}x^{1/2} + 6ax - 4a^{1/2}x^{3/2} + x^2] dx, \\
&= 1/6 \left[a^3 - 8/3a^3 + 3a^3 - 8/5a^3 + \frac{a^3}{3} \right] = \frac{a^3}{90}.
\end{aligned}$$

II. NOTES BY R. C. ARCHIBALD, Brown University.

The integral here arising is a particular case of one considered by Lejeune Dirichlet in 1839.¹

If $V = \int x^{a-1}y^{b-1}z^{c-1} \dots dx dy dz \dots$, considering all positive values of x, y, z, \dots such that

$$\left(\frac{x}{\alpha}\right)^p + \left(\frac{y}{\beta}\right)^q + \left(\frac{z}{\gamma}\right)^r + \dots < 1$$

the constants $a, b, c, \dots, p, q, r, \dots, \alpha, \beta, \gamma, \dots$ being also positive, then

$$V = \frac{\alpha^a \beta^b \gamma^c \dots}{pqr \dots} \frac{\Gamma\left(\frac{a}{p}\right) \Gamma\left(\frac{b}{q}\right) \Gamma\left(\frac{c}{r}\right) \dots}{\Gamma\left(1 + \frac{a}{p} + \frac{b}{q} + \frac{c}{r} + \dots\right)}.$$

For the surface $(x/a)^{1/n} + (y/b)^{1/n} + (z/c)^{1/n} = 1$, the volume in the first octant would be $abc(n!)^3/(3n)!$ When $n = 7/2$ we have a result given² in 1883

$$V = \frac{abc}{(2/7)^3} \cdot \frac{[\Gamma(7/2)]^3}{\Gamma(23/2)}.$$

The values, for $n = 3/2$ and $n = 1/2$ were given in Williamson, *Elementary Treatise on the Integral Calculus*, 6 ed., 1891, pp. 289-290. The result for the surface $(x/a)^2 + (y/b)^2 + (z/c)^4 = 1$, given in Todhunter, *A Treatise on the Integral Calculus*, 4 ed., 1874, p. 186, follows at once from the Dirichlet formula above.

2852 [1920, 377]. Proposed by D. H. RICHERT, Bethel College, Newton, Kans.

What is the radius of a cylinder inscribed in a right cone, radius of base $R = 5$ inches, and altitude $h = 18$ inches, the volume of the cylinder to be $1/n$ ($= 3/4$) that of the cone?

SOLUTION BY H. S. UHLER, Yale University.

Let V denote the volume of the cone, and let r, v , and z denote, respectively, the radius, the volume, and the altitude of the cylinder. Then

$$V = \frac{1}{3}\pi h R^2, \quad v = \pi z r^2, \quad \text{and} \quad v = \frac{1}{n} V;$$

hence,

$$z r^2 = \frac{h R^2}{3n}. \tag{1}$$

From the similar right triangles obtained by passing a plane through the common axis of the cone and cylinder we find

$$z = \frac{h(R-r)}{R}. \tag{2}$$

These two equations are homogeneous in h and z , and also in R and r ; therefore they determine the ratio of the altitudes and the ratio of the radii independently as functions of n alone.

Substituting from (2) for z in equation (1) we obtain

$$r^3 - R r^2 + R^3/(3n) = 0.$$

¹ *Comptes Rendus . . . de l'Académie des Sciences*, vol. 8, p. 156; also in *Journal de Mathématiques Pures et Appliquées*, vol. 4, p. 168.

² *Mathematical Questions with their solutions from the "Educational Times,"* vol. 38, p. 104.